antibodies -online.com









Overview

Quantity:	100 μL
Target:	Claudin 2 (CLDN2)
Binding Specificity:	C-Term
Reactivity:	Human, Mouse, Rat, Dog, Cow, Pig
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Claudin 2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF), Immunochromatography (IC)
	inimunocinomatography (ic)
Product Details	
Product Details Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of
Immunogen:	human Claudin 2.
Immunogen:	human Claudin 2.
Immunogen: Specificity:	human Claudin 2. Recognizes endogenous levels of Claudin 2 protein.
Immunogen: Specificity: Characteristics:	human Claudin 2. Recognizes endogenous levels of Claudin 2 protein. Rabbit polyclonal antibody to Claudin 2
Immunogen: Specificity: Characteristics: Purification:	human Claudin 2. Recognizes endogenous levels of Claudin 2 protein. Rabbit polyclonal antibody to Claudin 2

Target Details

Background:	Claudin-2, SP82
Gene ID:	9075, 12738
UniProt:	P57739, 088552
Pathways:	Hepatitis C


Application Details

Application Notes:	WB (1:500 - 1:1000), IH (1:100 - 1:200), IF/IC (1:100 - 1:500)
Restrictions:	For Research Use only

Handling


Format:	Liquid
Buffer:	Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.
Expiry Date:	12 months


Images



Western Blotting

Image 1. Western blot analysis of Claudin 2 expression in HeLa (A), A431 (B) whole cell lysates.





Immunohistochemistry

Image 2. Immunohistochemical analysis of Claudin 2 staining in human tonsil formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Immunofluorescence

Image 3. Immunofluorescent analysis of Claudin 2 staining in A431 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).