

Datasheet for ABIN7297528
anti-AIF antibody (N-Term)



[Go to Product page](#)

3 Images

Overview

Quantity:	100 µL
Target:	AIF (AIFM1)
Binding Specificity:	N-Term
Reactivity:	Human, Mouse, Rat, Sheep
Host:	Rabbit
Clonality:	Polyclonal
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF), Immunoprecipitation (IP), Immunochromatography (IC)

Product Details

Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human AIFM1.
Specificity:	Recognizes endogenous levels of AIFM1 protein.
Characteristics:	Rabbit polyclonal antibody to AIFM1
Purification:	The antibody was purified by immunogen affinity chromatography.

Target Details

Target:	AIF (AIFM1)
Alternative Name:	AIFM1 (AIFM1 Products)
Background:	AIF, PDCD8, Apoptosis-inducing factor 1, mitochondrial, Programmed cell death protein 8

Target Details

Gene ID:	9131, 26926, 83533
UniProt:	O95831 , Q9Z0X1 , Q9JM53
Pathways:	Apoptosis , Positive Regulation of Endopeptidase Activity , Cell Redox Homeostasis , Smooth Muscle Cell Migration , Warburg Effect


Application Details

Application Notes:	WB (1:500 - 1:1000), IH (1:100 - 1:200), IF/IC (1:100 - 1:500), IP (1:10 - 1:100)
Restrictions:	For Research Use only

Handling


Format:	Liquid
Buffer:	Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.
Expiry Date:	12 months

Images



Immunofluorescence


Image 1. Immunofluorescent analysis of AIFM1 staining in MCF7 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room



temperature in the dark. DAPI was used to stain the cell nuclei (blue).

Immunohistochemistry

Image 2. Immunohistochemical analysis of AIFM1 staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Western Blotting

Image 3. Western blot analysis of AIFM1 expression in MCF7 (A), Molt (B), Jurkat (C), mouse liver (D), rat heart (E) whole cell lysates.