antibodies -online.com





anti-NEMF antibody (C-Term)

2 Images



Go to Product page

\sim	
()\/\	rview
\cup	

Overview	
Quantity:	100 μL
Target:	NEMF (SDCCAG1)
Binding Specificity:	C-Term
Reactivity:	Human, Mouse, Monkey
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This NEMF antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC)
Product Details	
Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human NEMF.
Specificity:	Recognizes endogenous levels of NEMF protein.
Characteristics:	Rabbit polyclonal antibody to NEMF
Purification:	The antibody was purified by immunogen affinity chromatography.
Target Details	
Target:	NEMF (SDCCAG1)
Alternative Name:	NEMF (SDCCAG1 Products)
Background:	SDCCAG1, Nuclear export mediator factor NEMF, Antigen NY-CO-1, Serologically defined colon

Target Details

	cancer antigen 1
Gene ID:	9147, 66244
UniProt:	O60524, Q8CCP0


Application Details

Application Notes:	WB (1:500 - 1:1000), IH (1:100 - 1:200)
Restrictions:	For Research Use only

Handling


Format:	Liquid
Buffer:	Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.
Expiry Date:	12 months

Images



Immunohistochemistry

Image 1. Immunohistochemical analysis of NEMF staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Western Blotting

Image 2. Western blot analysis of NEMF expression in Jurkat (A) whole cell lysates.