

Datasheet for ABIN7297570  
**anti-DRAK2 antibody (Center)**[Go to Product page](#)

## 2 Images

## Overview

Quantity:	100 µL
Target:	DRAK2 (STK17B)
Binding Specificity:	Center
Reactivity:	Human, Mouse, Rat, Cow
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This DRAK2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC)

## Product Details

Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human STK17B.
Specificity:	Recognizes endogenous levels of STK17B protein.
Characteristics:	Rabbit polyclonal antibody to STK17B
Purification:	The antibody was purified by immunogen affinity chromatography.

## Target Details

Target:	DRAK2 (STK17B)
Alternative Name:	STK17B ( <a href="#">STK17B Products</a> )
Background:	DRAK2, Serine/threonine-protein kinase 17B, DAP kinase-related apoptosis-inducing protein

## Target Details

	kinase 2
Gene ID:	9262, 98267, 170904
UniProt:	<a href="#">O94768</a> , <a href="#">Q8BG48</a> , <a href="#">Q91XS8</a>


## Application Details

Application Notes:	WB (1:500 - 1:1000), IH (1:100 - 1:200)
Restrictions:	For Research Use only

## Handling


Format:	Liquid
Buffer:	Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.
Expiry Date:	12 months

## Images



### Immunohistochemistry

**Image 1.** Immunohistochemical analysis of STK17B staining in human lung cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Western Blotting

**Image 2.** Western blot analysis of STK17B expression in Jurkat (A), A673 (B) whole cell lysates.