antibodies -online.com





anti-GTF3C3 antibody (N-Term)

2 Images



Go to Product page

\sim					
	1//	r۱.	/ I	\triangle	٨

Overview	
Quantity:	100 μL
Target:	GTF3C3
Binding Specificity:	N-Term
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This GTF3C3 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC)
Product Details	
Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human GTF3C3.
Specificity:	Recognizes endogenous levels of GTF3C3 protein.
Characteristics:	Rabbit polyclonal antibody to GTF3C3
Purification:	The antibody was purified by immunogen affinity chromatography.
Target Details	
Target:	GTF3C3
Alternative Name:	GTF3C3 (GTF3C3 Products)
Background:	General transcription factor 3C polypeptide 3, Transcription factor IIIC 102 kDa subunit, TFIIIC

Target Details

	102 kDa subunit, TFIIIC102, Transcription factor IIIC subunit gamma, TF3C-gamma	
Gene ID:	9330	
UniProt:	Q9Y5Q9	


Application Details

Application Notes:	WB (1:500 - 1:1000), IH (1:100 - 1:200)
Restrictions:	For Research Use only

Handling


Format:	Liquid
Buffer:	Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.
Expiry Date:	12 months

Images



Immunohistochemistry

Image 1. Immunohistochemical analysis of GTF3C3 staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Western Blotting

Image 2. Western blot analysis of GTF3C3 expression in HeLa (A) whole cell lysates.