
antibodies - online.com







anti-EIF5B antibody (C-Term)





$C_0 + c$	Produc	+
	Promi	บ กลกย

Overview

Quantity:	100 μL
Target:	EIF5B
Binding Specificity:	C-Term
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This EIF5B antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP)

Product Details

Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human EIF5B.
Specificity:	Recognizes endogenous levels of EIF5B protein.
Characteristics:	Rabbit polyclonal antibody to EIF5B
Purification:	The antibody was purified by immunogen affinity chromatography.

Target Details

Target:	EIF5B
Alternative Name:	EIF5B (EIF5B Products)
Background:	IF2, KIAA0741, Eukaryotic translation initiation factor 5B, eIF-5B, Translation initiation factor IF-

Target Details

	2
Gene ID:	9669, 226982, 308306
UniProt:	O60841, Q05D44, B2GUV7


Application Details

Application Notes:	WB (1:500 - 1:1000), IH (1:100 - 1:200), IP (1:10 - 1:100)
Restrictions:	For Research Use only

Handling


Format:	Liquid
Buffer:	Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.
Expiry Date:	12 months

Images



Immunohistochemistry

Image 1. Immunohistochemical analysis of EIF5B staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Western Blotting

Image 2. Western blot analysis of EIF5B expression in A549 (A), Jurkat (B) whole cell lysates.