

Datasheet for ABIN7297708  
**anti-HDAC5 antibody (Center)**



[Go to Product page](#)

3 Images

## Overview

Quantity:	100 µL
Target:	HDAC5
Binding Specificity:	Center
Reactivity:	Human, Mouse, Rat, Cow, Pig
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This HDAC5 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF), Immunoprecipitation (IP), Immunochromatography (IC)

## Product Details

Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human Histone Deacetylase 5.
Specificity:	Recognizes endogenous levels of Histone Deacetylase 5 protein.
Characteristics:	Rabbit polyclonal antibody to Histone Deacetylase 5
Purification:	The antibody was purified by immunogen affinity chromatography.

## Target Details

Target:	HDAC5
Alternative Name:	Histone Deacetylase 5 ( <a href="#">HDAC5 Products</a> )

## Target Details


Background:	KIAA0600, Histone deacetylase 5, HD5, Antigen NY-CO-9
Gene ID:	10014
UniProt:	<a href="#">Q9UQL6</a> , <a href="#">Q9Z2V6</a>
Pathways:	<a href="#">Regulation of Muscle Cell Differentiation</a> , <a href="#">Skeletal Muscle Fiber Development</a> , <a href="#">Monocarboxylic Acid Catabolic Process</a>

## Application Details

Application Notes:	WB (1:500 - 1:1000), IH (1:100 - 1:200), IF/IC (1:100 - 1:500), IP (1:10 - 1:100)
Restrictions:	For Research Use only

## Handling

Format:	Liquid
Buffer:	Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.
Expiry Date:	12 months




### Western Blotting

**Image 1.** Western blot analysis of Histone Deacetylase 5 expression in MCF7 (A), HeLa (B), NIH3T3 (C), H9C2 (D) whole cell lysates.



### Immunofluorescence

**Image 2.** Immunofluorescent analysis of Histone Deacetylase 5 staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



### Immunohistochemistry

**Image 3.** Immunohistochemical analysis of Histone Deacetylase 5 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.