



Datasheet for ABIN7297796  
**anti-SYNCRIP antibody (Center)**



[Go to Product page](#)

2 Images

Overview

Quantity:	100 µL
Target:	SYNCRIP
Binding Specificity:	Center
Reactivity:	Human, Mouse, Rat, Dog, Cow, Rabbit, Zebrafish (Danio rerio), Pig
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SYNCRIP antibody is un-conjugated
Application:	Western Blotting (WB), Immunofluorescence (IF), Immunoprecipitation (IP), Immunochromatography (IC)

Product Details

Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human hnRNP Q.
Specificity:	Recognizes endogenous levels of hnRNP Q protein.
Characteristics:	Rabbit polyclonal antibody to hnRNP Q
Purification:	The antibody was purified by immunogen affinity chromatography.

Target Details

Target:	SYNCRIP
Alternative Name:	HnRNP Q ( <a href="#">SYNCRIP Products</a> )

## Target Details

---

Background:	HNRPQ, NSAP1, Heterogeneous nuclear ribonucleoprotein Q, hnRNP Q, Glycine- and tyrosine-rich RNA-binding protein, GRY-RBP, NS1-associated protein 1, Synaptotagmin-binding, cytoplasmic RNA-interacting protein
Gene ID:	10492, 56403, 363113
UniProt:	<a href="#">O60506</a> , <a href="#">Q7TMK9</a> , <a href="#">Q7TP47</a>

## Application Details


---

Application Notes:	WB (1:500 - 1:1000), IF/IC (1:100 - 1:500), IP (1:10 - 1:100)
Restrictions:	For Research Use only

## Handling


---

Format:	Liquid
Buffer:	Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.
Expiry Date:	12 months



### Immunofluorescence

**Image 1.** Immunofluorescent analysis of hnRNP Q staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



### Western Blotting

**Image 2.** Western blot analysis of hnRNP Q expression in HT1080 (A), HeLa (B), HuvEc (C) whole cell lysates.