antibodies -online.com





anti-USP19 antibody (Center)





Go to Product page

\sim					
	VE	٦r۱	/1/	$\triangle V$	٨
\cup	V	ر ار ا	יו ע	U V	V

Overview	
Quantity:	100 μL
Target:	USP19
Binding Specificity:	Center
Reactivity:	Human, Rat, Rabbit
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This USP19 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC)
Product Details	
Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human USP19.
Specificity:	Recognizes endogenous levels of USP19 protein.
Characteristics:	Rabbit polyclonal antibody to USP19
Purification:	The antibody was purified by immunogen affinity chromatography.
Target Details	
Target:	USP19
Alternative Name:	USP19 (USP19 Products)
Background:	KIAA0891, ZMYND9, Ubiquitin carboxyl-terminal hydrolase 19, Deubiquitinating enzyme 19,

Target Details

	Ubiquitin thioesterase 19, Ubiquitin-specific-processing protease 19, Zinc finger MYND domain-containing protein 9
Gene ID:	10869, 361190
UniProt:	O94966, Q6J1Y9


Application Details

Application Notes:	WB (1:500 - 1:1000), IH (1:100 - 1:200)
Restrictions:	For Research Use only

Handling


Format:	Liquid
Buffer:	Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.
Expiry Date:	12 months

Images



Immunohistochemistry

Image 1. Immunohistochemical analysis of USP19 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with



haematoxylin and mounted with DPX.

Western Blotting

Image 2. Western blot analysis of USP19 expression in Jurkat (A), H9C2 (B) whole cell lysates.