

Datasheet for ABIN7297856

anti-Peroxiredoxin 3 antibody (Center)**3** Images[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	Peroxiredoxin 3 (PRDX3)
Binding Specificity:	Center
Reactivity:	Human, Rat, Mouse, Monkey
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Peroxiredoxin 3 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF), Immunochromatography (IC)

Product Details

Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human Peroxiredoxin 3.
Specificity:	Recognizes endogenous levels of Peroxiredoxin 3 protein.
Characteristics:	Rabbit polyclonal antibody to Peroxiredoxin 3
Purification:	The antibody was purified by immunogen affinity chromatography.

Target Details

Target:	Peroxiredoxin 3 (PRDX3)
Alternative Name:	Peroxiredoxin 3 (PRDX3 Products)

Target Details


Background:	AOP1, Thioredoxin-dependent peroxide reductase, mitochondrial, Antioxidant protein 1, AOP-1, HBC189, Peroxiredoxin III, Prx-III, Peroxiredoxin-3, Protein MER5 homolog
Gene ID:	10935, 11757, 64371
UniProt:	P30048 , P20108 , Q9Z0V6
Pathways:	Ribonucleoside Biosynthetic Process , Methionine Biosynthetic Process

Application Details

Application Notes:	WB (1:500 - 1:1000), IH (1:100 - 1:200), IF/IC (1:100 - 1:500)
Restrictions:	For Research Use only


Handling

Format:	Liquid
Buffer:	Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.
Expiry Date:	12 months




Immunofluorescence

Image 1. Immunofluorescent analysis of Peroxiredoxin 3 staining in MCF7 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



Immunohistochemistry

Image 2. Immunohistochemical analysis of Peroxiredoxin 3 staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Western Blotting

Image 3. Western blot analysis of Peroxiredoxin 3 expression in HeLa (A), MCF7 (B) whole cell lysates.