

Datasheet for ABIN7298016
anti-BRMS1 antibody (Center)[Go to Product page](#)

2 Images

Overview

Quantity:	100 µL
Target:	BRMS1
Binding Specificity:	Center
Reactivity:	Human, Mouse, Rat, Monkey
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This BRMS1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC)

Product Details

Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human BRMS1.
Specificity:	Recognizes endogenous levels of BRMS1 protein.
Characteristics:	Rabbit polyclonal antibody to BRMS1
Purification:	The antibody was purified by immunogen affinity chromatography.

Target Details

Target:	BRMS1
Alternative Name:	BRMS1 (BRMS1 Products)
Background:	Breast cancer metastasis-suppressor 1

Target Details

Gene ID:	25855, 107392, 100911841
UniProt:	Q9HCU9 , Q99N20 , Q5M7T3


Application Details

Application Notes:	WB (1:500 - 1:1000), IH (1:100 - 1:200)
Restrictions:	For Research Use only

Handling


Format:	Liquid
Buffer:	Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.
Expiry Date:	12 months

Images



Immunohistochemistry

Image 1. Immunohistochemical analysis of BRMS1 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Western Blotting

Image 2. Western blot analysis of BRMS1 expression in HT29 (A), HeLa (B), MCF7 (C) whole cell lysates.