# antibodies -online.com





# anti-GORASP2 antibody (Center)

2 Images



Go to Product page

$\sim$					
()	VE	۲۱	/1	$\triangle$	Λ

Overview		
Quantity:	100 μL	
Target:	GORASP2	
Binding Specificity:	Center	
Reactivity:	Human, Mouse, Rat, Cow	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This GORASP2 antibody is un-conjugated	
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP)	
Product Details		
Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human GRASP55.	
Specificity:	Recognizes endogenous levels of GRASP55 protein.	
Characteristics:	Rabbit polyclonal antibody to GRASP55	
Purification:	The antibody was purified by immunogen affinity chromatography.	
Target Details		
Target:	GORASP2	
Alternative Name:	GRASP55 (GORASP2 Products)	
Background:	GOLPH6, Golgi reassembly-stacking protein 2, GRS2, Golgi phosphoprotein 6, GOLPH6, Golgi	

### **Target Details**

	reassembly-stacking protein of 55 kDa, GRASP55, p59	
Gene ID:	26003, 70231	
UniProt:	Q9H8Y8, Q99JX3, Q9R064	


### **Application Details**

Application Notes:	WB (1:500 - 1:1000), IH (1:100 - 1:200), IP (1:10 - 1:100)
Restrictions:	For Research Use only

### Handling


Format:	Liquid
Buffer:	Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.
Expiry Date:	12 months

#### **Images**



## **Immunohistochemistry**

Image 1. Immunohistochemical analysis of GRASP55 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



#### **Western Blotting**

Image 2. Western blot analysis of GRASP55 expression in HepG2 (A), Raji (B), LOVO (C) whole cell lysates.