# antibodies - online.com







## anti-Neurexophilin 1 antibody (Center)





<i>ا</i> ۱	1	-	M	/1	0	
u	1//	$\vdash$	1 \	/ I	-	\/\/
$\sim$	v	$\sim$	r٧		$\sim$	v v

Overview	
Quantity:	100 μL
Target:	Neurexophilin 1 (NXPH1)
Binding Specificity:	Center
Reactivity:	Human, Dog, Cow, Monkey
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Neurexophilin 1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC)
Product Details	
Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human NXPH1.
Specificity:	Recognizes endogenous levels of NXPH1 protein.
Characteristics:	Rabbit polyclonal antibody to NXPH1
Purification:	The antibody was purified by immunogen affinity chromatography.
Target Details	
Target:	Neurexophilin 1 (NXPH1)
Alternative Name:	NXPH1 (NXPH1 Products)
Background:	NPH1, Neurexophilin-1

### **Target Details**

Gene ID:	30010
UniProt:	P58417

## **Application Details**

Application Notes:	WB (1:500 - 1:1000), IH (1:100 - 1:200)	
Restrictions:	For Research Use only	

## Handling

**Images** 

Format:	Liquid
Buffer:	Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.
Expiry Date:	12 months




kDa

60

50 40 30

#### **Western Blotting**

**Image 1.** Western blot analysis of NXPH1 expression in A549 (A) whole cell lysates.



#### **Immunohistochemistry**

Image 2. Immunohistochemical analysis of NXPH1 staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.