
antibodies - online.com







anti-CDC40 antibody (Center)





\sim				
	11/6	٦r١	/10	۱۸.

Overview		
Quantity:	100 μL	
Target:	CDC40	
Binding Specificity:	Center	
Reactivity:	Mouse, Rat, Human, Cow	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This CDC40 antibody is un-conjugated	
Application:	Western Blotting (WB), Immunoprecipitation (IP), Immunohistochemistry (IHC)	
Product Details		
Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human CDC40.	
Specificity:	Recognizes endogenous levels of CDC40 protein.	
Characteristics:	Rabbit polyclonal antibody to CDC40	
Purification:	The antibody was purified by immunogen affinity chromatography.	
Target Details		
Target:	CDC40	
Alternative Name:	CDC40 (CDC40 Products)	
Background:	EHB3, PRP17, PRPF17, Pre-mRNA-processing factor 17, Cell division cycle 40 homolog, EH-	

Target Details

	binding protein 3, Ehb3, PRP17 homolog, hPRP17
Gene ID:	51362, 71713
UniProt:	O60508, Q9DC48
Pathways:	Ribonucleoprotein Complex Subunit Organization


Application Details

Application Notes:	WB (1:500 - 1:1000), IH (1:100 - 1:200), IP (1:10 - 1:100)
Restrictions:	For Research Use only

Handling


Format:	Liquid
Buffer:	Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.
Expiry Date:	12 months

Images



Western Blotting

Image 1. Western blot analysis of CDC40 expression in K562 (A), HuvEC (B) whole cell lysates.



Immunohistochemistry

Image 2. Immunohistochemical analysis of CDC40 staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.