
# antibodies -online.com







# anti-TOLLIP antibody (Center)

**Images** 

Overview

Characteristics:

**Target Details** 

Purification:



Quantity:	100 μL
Target:	TOLLIP
Binding Specificity:	Center
Reactivity:	Human, Monkey
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TOLLIP antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP)
Product Details	
Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human
	TOLLIP.
Specificity:	Recognizes endogenous levels of TOLLIP protein.

Target:	TOLLIP	

Rabbit polyclonal antibody to TOLLIP

Alternative Name: TOLLIP (TOLLIP Products)

Background: Toll-interacting protein

The antibody was purified by immunogen affinity chromatography.

## **Target Details**

Gene ID:	54472
UniProt:	Q9H0E2


# **Application Details**

Application Notes:	WB (1:500 - 1:1000), IH (1:100 - 1:200), IP (1:10 - 1:100)
Restrictions:	For Research Use only

## Handling


Format:	Liquid
Buffer:	Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.
Expiry Date:	12 months

#### **Images**



#### Immunohistochemistry

Image 1. Immunohistochemical analysis of TOLLIP staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



#### **Western Blotting**

Image 2. Western blot analysis of TOLLIP expression in Jurkat (A), LNCaP (B), HepG2 (C) whole cell lysates.