



[Go to Product page](#)

Datasheet for ABIN7298276

anti-Cyclin L1 antibody (C-Term)

2 Images

Overview

Quantity:	100 µL
Target:	Cyclin L1 (CCNL1)
Binding Specificity:	C-Term
Reactivity:	Human, Mouse, Rat, Dog, Cow, Pig
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Cyclin L1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC)

Product Details

Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human Cyclin L1.
Specificity:	Recognizes endogenous levels of Cyclin L1 protein.
Characteristics:	Rabbit polyclonal antibody to Cyclin L1
Purification:	The antibody was purified by immunogen affinity chromatography.

Target Details

Target:	Cyclin L1 (CCNL1)
Alternative Name:	Cyclin L1 (CCNL1 Products)
Background:	Cyclin-L1, Cyclin-L

Target Details

Gene ID: 57018, 56706, 100909712

UniProt: [Q9UK58](#), [Q52KE7](#), [Q9R1Q2](#)

Application Details

Application Notes: WB (1:500 - 1:1000), IH (1:100 - 1:200)

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.

Preservative: Sodium azide


Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.


Expiry Date: 12 months

Images



Western Blotting

Image 1. Western blot analysis of Cyclin L1 expression in HL60 (A), NIH3T3 (B), Jurkat (C), THP1 (D), K562 (E) whole cell lysates.



Immunohistochemistry

Image 2. Immunohistochemical analysis of Cyclin L1 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.