

Datasheet for ABIN7298284
anti-PAK7 antibody (C-Term)[Go to Product page](#)

2 Images

Overview

Quantity:	100 µL
Target:	PAK7
Binding Specificity:	C-Term
Reactivity:	Human, Mouse, Rat, Monkey
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PAK7 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC)

Product Details

Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human PAK7.
Specificity:	Recognizes endogenous levels of PAK7 protein.
Characteristics:	Rabbit polyclonal antibody to PAK7
Purification:	The antibody was purified by immunogen affinity chromatography.

Target Details

Target:	PAK7
Alternative Name:	PAK7 (PAK7 Products)
Background:	KIAA1264, PAK5, Serine/threonine-protein kinase PAK 7, p21-activated kinase 5, PAK-5, p21-

Target Details

activated kinase 7, PAK-7

Gene ID: 57144, 241656, 311450

UniProt: [Q9P286](#), [Q8C015](#), [D4A280](#)

Application Details

Application Notes: WB (1:500 - 1:1000), IH (1:100 - 1:200)

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.

Preservative: Sodium azide


Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.


Expiry Date: 12 months

Images



Immunohistochemistry

Image 1. Immunohistochemical analysis of PAK7 staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Western Blotting

Image 2. Western blot analysis of PAK7 expression in SKNSH (A), K562 (B), COS7 (C) whole cell lysates.