

Datasheet for ABIN7298416
anti-CSRNP2 antibody (Center)[Go to Product page](#)

2 Images

Overview

Quantity:	100 µL
Target:	CSRNP2
Binding Specificity:	Center
Reactivity:	Human, Mouse, Rat, Monkey
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CSRNP2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP)

Product Details

Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human CSRNP2.
Specificity:	Recognizes endogenous levels of CSRNP2 protein.
Characteristics:	Rabbit polyclonal antibody to CSRNP2
Purification:	The antibody was purified by immunogen affinity chromatography.

Target Details

Target:	CSRNP2
Alternative Name:	CSRNP2 (CSRNP2 Products)
Background:	C12orf22, FAM130A1, TAIP12, Cysteine/serine-rich nuclear protein 2, CSRNP-2, Protein

Target Details

FAM130A1, TGF-beta-induced apoptosis protein 12, TAIP-12

Gene ID: 81566, 207785

UniProt: [Q9H175](#), [Q8BGQ2](#)

Application Details

Application Notes: WB (1:500 - 1:1000), IH (1:100 - 1:200), IP (1:10 - 1:100)

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.

Preservative: Sodium azide


Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.


Expiry Date: 12 months

Images



Immunohistochemistry

Image 1. Immunohistochemical analysis of CSRNP2 staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Western Blotting

Image 2. Western blot analysis of CSRNP2 expression in HEK293T (A), HepG2 (B), HuvEC (C) whole cell lysates.