



[Go to Product page](#)

Datasheet for ABIN7298466
anti-RELT antibody (C-Term)

2 Images

Overview

Quantity:	100 µL
Target:	RELT
Binding Specificity:	C-Term
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This RELT antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP)

Product Details

Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human RELT.
Specificity:	Recognizes endogenous levels of RELT protein.
Characteristics:	Rabbit polyclonal antibody to RELT
Purification:	The antibody was purified by immunogen affinity chromatography.

Target Details

Target:	RELT
Alternative Name:	RELT (RELT Products)
Background:	TNFRSF19L, Tumor necrosis factor receptor superfamily member 19L, Receptor expressed in

Target Details

lymphoid tissues

Gene ID: 84957, 320100

UniProt: [Q969Z4](#), [Q8BX43](#)

Application Details

Application Notes: WB (1:500 - 1:1000), IH (1:100 - 1:200), IP (1:10 - 1:100)

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.

Preservative: Sodium azide


Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.


Expiry Date: 12 months

Images



Western Blotting

Image 1. Western blot analysis of RELT expression in HeLa (A), T47D (B), COLO205 (C) whole cell lysates.



Immunohistochemistry

Image 2. Immunohistochemical analysis of RELT staining in human lymph node formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.