antibodies -online.com





anti-TP53INP1 antibody (C-Term)

2 Images



Go to Product page

\sim				
	11/6	٦r١	/10	۱۸.

Overview		
Quantity:	100 μL	
Target:	TP53INP1	
Binding Specificity:	C-Term	
Reactivity:	Human, Monkey	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This TP53INP1 antibody is un-conjugated	
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP)	
Product Details		
Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human TP53INP1.	
Specificity:	Recognizes endogenous levels of TP53INP1 protein.	
Characteristics:	Rabbit polyclonal antibody to TP53INP1	
Purification:	The antibody was purified by immunogen affinity chromatography.	
Target Details		
Target:	TP53INP1	
Alternative Name:	TP53INP1 (TP53INP1 Products)	
Background:	P53DINP1, SIP, Tumor protein p53-inducible nuclear protein 1, Stress-induced protein, p53-	

Target Details

	dependent damage-inducible nuclear protein 1, p53DINP1
Gene ID:	94241
UniProt:	Q96A56


Application Details

Application Notes:	WB (1:500 - 1:1000), IH (1:100 - 1:200), IP (1:10 - 1:100)
Restrictions:	For Research Use only

Handling


Format:	Liquid
Buffer:	Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.
Expiry Date:	12 months

Images



Immunohistochemistry

Image 1. Immunohistochemical analysis of TP53INP1 staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Western Blotting

Image 2. Western blot analysis of TP53INP1 expression in HeLa (A), HepG2 (B) whole cell lysates.