

[Go to Product page](#)

## Datasheet for ABIN729900 **anti-MAFA antibody (Biotin)**

### Overview

Quantity:	100 µL
Target:	MAFA
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MAFA antibody is conjugated to Biotin
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

### Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Mafa
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Purified by Protein A.

### Target Details

Target:	MAFA
Alternative Name:	Mafa ( <a href="#">MAFA Products</a> )
Background:	Synonyms: Mafa homolog, V-maf musculoaponeurotic fibrosarcoma oncogene homolog A, Pancreatic beta-cell-specific transcriptional activator, avian V-maf musculoaponeurotic fibrosarcoma oncogene homolog A, MAFA_MOUSE. Background: Insulin gene expression is regulated by several islet-enriched transcription factors.

## Target Details

However, MAFA is the only beta cell-specific activator. MAFA selectively induces endogenous insulin transcription in non-beta cells. MAFA was also first detected in the insulin-producing cells formed during the second and predominant phase of beta cell differentiation, and absent in the few insulin-positive cells found in Nkx6.1(-/-) pancreata, which lack the majority of second-phase beta cells. These results demonstrate that MAFA is a potent insulin activator that is likely to function downstream of Nkx6.1 during islet insulin-producing cell development.

Gene ID:	389692
Pathways:	<a href="#">Hormone Transport</a>

## Application Details

Application Notes:	WB: (1:100-1000), IHC-P: (1:100-500) Optimal working dilution should be determined by the investigator.
Restrictions:	For Research Use only

## Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months