



[Go to Product page](#)

Datasheet for ABIN7299114
anti-SNRPE antibody (C-Term)

2 Images

Overview

Quantity:	100 µL
Target:	SNRPE
Binding Specificity:	C-Term
Reactivity:	Human, Mouse, Rat, Chicken, Zebrafish (Danio rerio)
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SNRPE antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC)

Product Details

Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human B-RAF.
Specificity:	Recognizes endogenous levels of B-RAF protein.
Characteristics:	Rabbit polyclonal antibody to B-RAF
Purification:	The antibody was purified by immunogen affinity chromatography.

Target Details

Target:	SNRPE
Alternative Name:	B-RAF (SNRPE Products)
Background:	BRAF1, RAFB1, Serine/threonine-protein kinase B-raf, Proto-oncogene B-Raf, p94, v-Raf murine

Target Details

sarcoma viral oncogene homolog B1

Gene ID: 673, 109880

UniProt: [P15056](#), [P28028](#)

Pathways: [Ribonucleoprotein Complex Subunit Organization](#), [Hepatitis C](#)

Application Details

Application Notes: WB (1:500 - 1:1000), IH (1:100 - 1:200)

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.

Preservative: Sodium azide


Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.

Expiry Date: 12 months


Images



Immunohistochemistry

Image 1. Immunohistochemical analysis of B-RAF staining in human prostate cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with

haematoxylin and mounted with DPX.



Western Blotting

Image 2. Western blot analysis of B-RAF expression in HL60 (A), HeLa (B), NIH3T3 (C) whole cell lysates.