

Datasheet for ABIN7299232

anti-eIF4EBP1 antibody (N-Term, pSer37)



[Go to Product page](#)

2 Images

Overview

Quantity:	100 µL
Target:	eIF4EBP1 (EIF4EBP1)
Binding Specificity:	N-Term, pSer37
Reactivity:	Human, Mouse, Rat, Cow, Pig
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This eIF4EBP1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC)

Product Details

Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human 4EBP1.
Specificity:	Recognizes endogenous levels of 4EBP1 (pT37) protein.
Characteristics:	Rabbit polyclonal antibody to 4EBP1 (pT37)
Purification:	The antibody was purified by immunogen affinity chromatography.

Target Details

Target:	eIF4EBP1 (EIF4EBP1)
Alternative Name:	4EBP1 (EIF4EBP1 Products)
Background:	Eukaryotic translation initiation factor 4E-binding protein 1, 4E-BP1, eIF4E-binding protein 1,

Target Details


	Phosphorylated heat- and acid-stable protein regulated by insulin 1, PHAS-I
Gene ID:	1978, 13685, 116636
UniProt:	Q13541 , Q60876 , Q62622
Pathways:	MAPK Signaling , PI3K-Akt Signaling , RTK Signaling , AMPK Signaling , Regulation of Cell Size , BCR Signaling

Application Details

Application Notes:	WB (1:500 - 1:1000), IH (1:100 - 1:200)
Restrictions:	For Research Use only


Handling

Format:	Liquid
Buffer:	Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.
Expiry Date:	12 months



Immunohistochemistry

Image 1. Immunohistochemical analysis of 4EBP1 (pT37) staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Western Blotting

Image 2. Western blot analysis of 4EBP1 (pT37) expression in HEK293T insulin-treated (A), NIH3T3 insulin-treated (B), PC12 insulin-treated (C) whole cell lysates.