



Datasheet for ABIN7299526
anti-ZNF174 antibody (Center)



[Go to Product page](#)

3 Images

Overview

Quantity:	100 µL
Target:	ZNF174
Binding Specificity:	Center
Reactivity:	Human, Rat, Monkey, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ZNF174 antibody is un-conjugated
Application:	Western Blotting (WB), Immunofluorescence (IF), Immunohistochemistry (IHC), Immunochromatography (IC)

Product Details

Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human ZNF174.
Specificity:	Recognizes endogenous levels of ZNF174 protein.
Characteristics:	Rabbit polyclonal antibody to ZNF174
Purification:	The antibody was purified by immunogen affinity chromatography.

Target Details

Target:	ZNF174
Alternative Name:	ZNF174 (ZNF174 Products)

Target Details

Background:	ZSCAN8, Zinc finger protein 174, AW-1, Zinc finger and SCAN domain-containing protein 8
Gene ID:	7727
UniProt:	Q15697


Application Details

Application Notes:	WB (1:500 - 1:1000), IH (1:100 - 1:200), IF/IC (1:100 - 1:500)
Restrictions:	For Research Use only

Handling


Format:	Liquid
Buffer:	Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.
Expiry Date:	12 months

Images



Immunofluorescence


Image 1. Immunofluorescent analysis of ZNF174 staining in Jurkat cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell



nuclei (blue).

Immunohistochemistry

Image 2. Immunohistochemical analysis of ZNF174 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Western Blotting

Image 3. Western blot analysis of ZNF174 expression in Jurkat (A), Raw264.7 (B), H9C2 (C) whole cell lysates.