



[Go to Product page](#)

Datasheet for ABIN7299542

## anti-Cyclin A1 antibody (C-Term)

### 3 Images

#### Overview

Quantity:	100 µL
Target:	Cyclin A1 (CCNA1)
Binding Specificity:	C-Term
Reactivity:	Human, Mouse, Rat, Cow, Rabbit, Chicken, Dog, Pig
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Cyclin A1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF), Immunochromatography (IC)

#### Product Details

Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human Cyclin A1.
Specificity:	Recognizes endogenous levels of Cyclin A1 protein.
Characteristics:	Rabbit polyclonal antibody to Cyclin A1
Purification:	The antibody was purified by immunogen affinity chromatography.

#### Target Details

Target:	Cyclin A1 (CCNA1)
Alternative Name:	Cyclin A1 ( <a href="#">CCNA1 Products</a> )

## Target Details

---

Background:	Cyclin-A1
Gene ID:	8900, 12427, 295052
UniProt:	<a href="#">P78396</a> , <a href="#">Q61456</a> , <a href="#">Q6AY13</a>
Pathways:	<a href="#">Apoptosis</a> , <a href="#">Cell Division Cycle</a> , <a href="#">AMPK Signaling</a> , <a href="#">Mitotic G1-G1/S Phases</a> , <a href="#">DNA Replication</a> , <a href="#">M Phase</a> , <a href="#">Synthesis of DNA</a>

## Application Details


---

Application Notes:	WB (1:500 - 1:1000), IH (1:100 - 1:200), IF/IC (1:100 - 1:500)
Restrictions:	For Research Use only

## Handling


---

Format:	Liquid
Buffer:	Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.
Expiry Date:	12 months




### Immunofluorescence

**Image 1.** Immunofluorescent analysis of Cyclin A1 staining in MCF7 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



### Immunohistochemistry

**Image 2.** Immunohistochemical analysis of Cyclin A1 staining in human testis formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



### Western Blotting

**Image 3.** Western blot analysis of Cyclin A1 expression in MCF7 (A), NIH3T3 (B) whole cell lysates.