# antibodies - online.com







## anti-CDKL2 antibody (Center)





$\sim$				
	11/6	٦r١	/10	۱۸.

Overview		
Quantity:	100 μL	
Target:	CDKL2	
Binding Specificity:	Center	
Reactivity:	Human, Monkey	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This CDKL2 antibody is un-conjugated	
Application:	Western Blotting (WB), Immunohistochemistry (IHC)	
Product Details		
Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human CDKL2.	
Specificity:	Recognizes endogenous levels of CDKL2 protein.	
Characteristics:	Rabbit polyclonal antibody to CDKL2	
Purification:	The antibody was purified by immunogen affinity chromatography.	
Target Details		
Target:	CDKL2	
Alternative Name:	CDKL2 (CDKL2 Products)	
Background:	Cyclin-dependent kinase-like 2, Protein kinase p56 KKIAMRE, Serine/threonine-protein kinase	

#### **Target Details**

	KKIAMRE
Gene ID:	8999
UniProt:	Q92772


## **Application Details**

Application Notes:	WB (1:500 - 1:1000), IH (1:100 - 1:200)
Restrictions:	For Research Use only

## Handling


Format:	Liquid
Buffer:	Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.
Expiry Date:	12 months

#### **Images**



## **Western Blotting**

Image 1. Western blot analysis of CDKL2 expression in SHSYSY (A) whole cell lysates.



#### **Immunohistochemistry**

Image 2. Immunohistochemical analysis of CDKL2 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.