

Datasheet for ABIN7299656

anti-HDAC8 antibody (N-Term, pSer39)[Go to Product page](#)**3** Images

Overview

Quantity:	100 µL
Target:	HDAC8
Binding Specificity:	N-Term, pSer39
Reactivity:	Human, Mouse, Rat, Monkey
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This HDAC8 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF), Immunochromatography (IC)

Product Details

Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human Histone Deacetylase 8.
Specificity:	Recognizes endogenous levels of Histone Deacetylase 8 (pS39) protein.
Characteristics:	Rabbit polyclonal antibody to Histone Deacetylase 8 (pS39)
Purification:	The antibody was purified by immunogen affinity chromatography.

Target Details

Target:	HDAC8
Alternative Name:	Histone Deacetylase 8 (HDAC8 Products)

Target Details

Background:	HDACL1, Histone deacetylase 8, HD8
Gene ID:	55869, 70315, 100911968
UniProt:	Q9BY41 , Q8VH37 , B1WC68
Pathways:	Cellular Glucan Metabolic Process


Application Details

Application Notes:	WB (1:500 - 1:1000), IH (1:100 - 1:200), IF/IC (1:100 - 1:500)
Restrictions:	For Research Use only

Handling


Format:	Liquid
Buffer:	Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.
Expiry Date:	12 months

Images



Immunofluorescence


Image 1. Immunofluorescent analysis of Histone Deacetylase 8 (pS39) staining in A549 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary



antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).

Immunohistochemistry

Image 2. Immunohistochemical analysis of Histone Deacetylase 8 (pS39) staining in human lung cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Western Blotting

Image 3. Western blot analysis of Histone Deacetylase 8 (pS39) expression in A549 (A), NIH3T3 (B), rat kidney (C) whole cell lysates.