# antibodies - online.com







# anti-BAIAP2L2 antibody (Center)

**Images** 



$\sim$				
	11/6	٦r١	/10	۱۸.

Quantity:	100 μL	
Target:	BAIAP2L2	
Binding Specificity:	Center	
Reactivity:	Human, Mouse, Cow	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This BAIAP2L2 antibody is un-conjugated	
Application:	Western Blotting (WB), Immunohistochemistry (IHC)	
Product Details		
Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human BAIAP2L2.	
Specificity:	Recognizes endogenous levels of BAIAP2L2 protein.	
Characteristics:	Rabbit polyclonal antibody to BAIAP2L2	
Purification:	The antibody was purified by immunogen affinity chromatography.	
Target Details		
Target:	BAIAP2L2	
Alternative Name:	BAIAP2L2 (BAIAP2L2 Products)	
Background:	Brain-specific angiogenesis inhibitor 1-associated protein 2-like protein 2, BAI1-associated	

### **Target Details**

	protein 2-like protein 2, Planar intestinal- and kidney-specific BAR domain protein, Pinkbar
Gene ID:	80115, 207495
UniProt:	Q6UXY1, Q80Y61


### **Application Details**

Application Notes:	WB (1:500 - 1:1000), IH (1:100 - 1:200)
Restrictions:	For Research Use only

### Handling


Format:	Liquid
Buffer:	Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.
Expiry Date:	12 months

#### **Images**



## **Immunohistochemistry**

Image 1. Immunohistochemical analysis of BAIAP2L2 staining in human lung cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



#### **Western Blotting**

Image 2. Western blot analysis of BAIAP2L2 expression in HeLa (A), HL60 (B), NIH3T3 (C) whole cell lysates.