



[Go to Product page](#)

Datasheet for ABIN7299710
anti-USP32 antibody (Center)

2 Images

Overview

Quantity:	100 µL
Target:	USP32
Binding Specificity:	Center
Reactivity:	Human, Mouse, Dog, Rabbit, Sheep
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This USP32 antibody is un-conjugated
Application:	Immunohistochemistry (IHC), Western Blotting (WB)

Product Details

Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human USP32.
Specificity:	Recognizes endogenous levels of USP32 protein.
Characteristics:	Rabbit polyclonal antibody to USP32
Purification:	The antibody was purified by immunogen affinity chromatography.

Target Details

Target:	USP32
Alternative Name:	USP32 (USP32 Products)
Background:	USP10, Ubiquitin carboxyl-terminal hydrolase 32, Deubiquitinating enzyme 32, Renal carcinoma

Target Details

antigen NY-REN-60, Ubiquitin thioesterase 32, Ubiquitin-specific-processing protease 32

Gene ID: 84669

UniProt: [Q8NFA0](#)

Application Details

Application Notes: WB (1:500 - 1:1000), IH (1:100 - 1:200)

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.

Preservative: Sodium azide


Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.


Expiry Date: 12 months

Images



Immunohistochemistry

Image 1. Immunohistochemical analysis of USP32 staining in human lung cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Western Blotting

Image 2. Western blot analysis of USP32 expression in HeLa (A), NIH3T3 (B) whole cell lysates.