



Datasheet for ABIN7299746
anti-ANKRD20A3 antibody (Center)



[Go to Product page](#)

2 Images

Overview

Quantity:	100 µL
Target:	ANKRD20A3
Binding Specificity:	Center
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ANKRD20A3 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC)

Product Details

Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human ANKRD20A3.
Specificity:	Recognizes endogenous levels of ANKRD20A3 protein.
Characteristics:	Rabbit polyclonal antibody to ANKRD20A3
Purification:	The antibody was purified by immunogen affinity chromatography.

Target Details

Target:	ANKRD20A3
Alternative Name:	ANKRD20A3 (ANKRD20A3 Products)
Background:	Ankyrin repeat domain-containing protein 20A3

Target Details

Gene ID: 441425

UniProt: [Q5VUR7](#)

Application Details

Application Notes: WB (1:500 - 1:1000), IH (1:100 - 1:200)

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.

Preservative: Sodium azide


Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.


Expiry Date: 12 months

Images



Western Blotting

Image 1. Western blot analysis of ANKRD20A3 expression in Jurkat (A), K562 (B) whole cell lysates.



Immunohistochemistry

Image 2. Immunohistochemical analysis of ANKRD20A3 staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.