



[Go to Product page](#)

Datasheet for ABIN7300428  
**anti-BMX antibody (N-Term)**

3 Images

### Overview

|                      |   |
|----------------------|---|
| Quantity:            | 100 µL  |
| Target:              | BMX   |
| Binding Specificity: | N-Term  |
| Reactivity:          | Human, Mouse, Rat   |
| Host:                | Rabbit  |
| Clonality:           | Polyclonal  |
| Conjugate:           | This BMX antibody is un-conjugated  |
| Application:         | Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF), Immunoprecipitation (IP), Immunochromatography (IC) |

### Product Details

|                  |   |
|------------------|---|
| Immunogen:       | KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human BMX. |
| Specificity:     | Recognizes endogenous levels of BMX protein.  |
| Characteristics: | Rabbit polyclonal antibody to BMX   |
| Purification:    | The antibody was purified by immunogen affinity chromatography.                                 |

### Target Details

|                   |                                      |
|-------------------|--------------------------------------|
| Target:           | BMX                                  |
| Alternative Name: | BMX ( <a href="#">BMX Products</a> ) |

## Target Details

|             |   |
|-------------|---|
| Background: | Cytoplasmic tyrosine-protein kinase BMX, Bone marrow tyrosine kinase gene in chromosome X protein, Epithelial and endothelial tyrosine kinase, ETK, NTK38 |
| Gene ID:    | 660, 12169  |
| UniProt:    | <a href="#">P51813</a> , <a href="#">P97504</a>   |


## Application Details

|                    |   |
|--------------------|---|
| Application Notes: | WB (1:500 - 1:1000), IH (1:100 - 1:200), IF/IC (1:100 - 1:500), IP (1:10 - 1:100) |
| Restrictions:      | For Research Use only   |

## Handling


|                    |  |
|--------------------|--|
| Format:            | Liquid   |
| Buffer:            | Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.          |
| Preservative:      | Sodium azide   |
| Precaution of Use: | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Storage:           | -20 °C   |
| Storage Comment:   | Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.                       |
| Expiry Date:       | 12 months  |

## Images




### Western Blotting

**Image 1.** Western blot analysis of BMX expression in mouse brain (A) whole cell lysates.



### Immunofluorescence

**Image 2.** Immunofluorescent analysis of BMX staining in A549 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



### Immunohistochemistry

**Image 3.** Immunohistochemical analysis of BMX staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX. w