# antibodies - online.com







# anti-JMY antibody (C-Term)

**Images** 



( )	1 /	$\sim$	KI /	110	Νę
	1//	$\vdash$	I \/	1 ←	٠// ٢

Overview		
Quantity:	100 μL	
Target:	JMY	
Binding Specificity:	C-Term	
Reactivity:	Human, Mouse, Cow, Dog, Chicken, Pig	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This JMY antibody is un-conjugated	
Application:	Western Blotting (WB), Immunohistochemistry (IHC)	
Product Details		
Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human JMY.	
Specificity:	Recognizes endogenous levels of JMY protein.	
Characteristics:	Rabbit polyclonal antibody to JMY	
Purification:	The antibody was purified by immunogen affinity chromatography.	
Target Details		
Target:	JMY	
Alternative Name:	JMY (JMY Products)	
Background:	Junction-mediating and -regulatory protein	

#### **Target Details**

Gene ID:	133746, 57748	
UniProt:	Q8N9B5, Q9QXM1	
Pathways:	Regulation of Actin Filament Polymerization	


# **Application Details**

Application Notes:	WB (1:500 - 1:1000), IH (1:100 - 1:200)
Restrictions:	For Research Use only

## Handling


Format:	Liquid
Buffer:	Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.
Expiry Date:	12 months

#### **Images**



## **Immunohistochemistry**

Image 1. Immunohistochemical analysis of JMY staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugad compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



#### **Western Blotting**

Image 2. Western blot analysis of JMY expression in Jurkat (A), MCF7 (B) whole cell lysates.