

Datasheet for ABIN7301024
anti-MYF5 antibody (Center)

3 Images

[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	MYF5
Binding Specificity:	Center
Reactivity:	Human, Monkey
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MYF5 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF), Immunochromatography (IC)

Product Details

Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human MYF5.
Specificity:	Recognizes endogenous levels of MYF5 protein.
Characteristics:	Rabbit polyclonal antibody to MYF5
Purification:	The antibody was purified by immunogen affinity chromatography.

Target Details

Target:	MYF5
Alternative Name:	MYF5 (MYF5 Products)

Target Details

Background:	BHLHC2, Myogenic factor 5, Myf-5, Class C basic helix-loop-helix protein 2, bHLHc2
Gene ID:	4617, 17877
UniProt:	P13349 , P24699
Pathways:	Regulation of Muscle Cell Differentiation , Skeletal Muscle Fiber Development


Application Details

Application Notes:	WB (1:500 - 1:1000), IH (1:100 - 1:200), IF/IC (1:100 - 1:500)
Restrictions:	For Research Use only

Handling

Format:	Liquid
Buffer:	Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.
Expiry Date:	12 months


Images



Immunofluorescence

Image 1. Immunofluorescent analysis of MYF5 staining in Jurkat cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room

temperature in the dark. DAPI was used to stain the cell nuclei (blue).



Immunohistochemistry

Image 2. Immunohistochemical analysis of MYF5 staining in human lung cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Western Blotting

Image 3. Western blot analysis of MYF5 expression in Jurkat (A) whole cell lysates.

