

Datasheet for ABIN7301066
anti-NOM1 antibody (C-Term)[Go to Product page](#)

2 Images

Overview

| | |
|----------------------|---|
| Quantity: | 100 µL |
| Target: | NOM1 |
| Binding Specificity: | C-Term |
| Reactivity: | Human, Rat |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This NOM1 antibody is un-conjugated |
| Application: | Immunohistochemistry (IHC), Western Blotting (WB) |

Product Details

| | |
|------------------|--|
| Immunogen: | KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human NOM1. |
| Specificity: | Recognizes endogenous levels of NOM1 protein. |
| Characteristics: | Rabbit polyclonal antibody to NOM1 |
| Purification: | The antibody was purified by immunogen affinity chromatography. |

Target Details

| | |
|-------------------|---|
| Target: | NOM1 |
| Alternative Name: | NOM1 (NOM1 Products) |
| Background: | C7orf3, Nucleolar MIF4G domain-containing protein 1, SGD1 homolog |

Target Details

| | |
|----------|---|
| Gene ID: | 64434, 433864 |
| UniProt: | Q5C9Z4 , Q3UFM5 |


Application Details

| | |
|--------------------|---|
| Application Notes: | WB (1:500 - 1:1000), IH (1:100 - 1:200) |
| Restrictions: | For Research Use only |

Handling


| | |
|--------------------|--|
| Format: | Liquid |
| Buffer: | Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide. |
| Preservative: | Sodium azide |
| Precaution of Use: | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Storage: | -20 °C |
| Storage Comment: | Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles. |
| Expiry Date: | 12 months |

Images



Western Blotting

Image 1. Western blot analysis of NOM1 expression in HEK293T (A), H9C2 (B) whole cell lysates.



Immunohistochemistry

Image 2. Immunohistochemical analysis of NOM1 staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.