

Datasheet for ABIN7301122  
**anti-PIGF antibody (C-Term)**

## 2 Images

[Go to Product page](#)

## Overview

Quantity:	100 µL
Target:	PIGF
Binding Specificity:	C-Term
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PIGF antibody is un-conjugated
Application:	Immunohistochemistry (IHC), Western Blotting (WB)

## Product Details

Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human PIGF.
Specificity:	Recognizes endogenous levels of PIGF protein.
Characteristics:	Rabbit polyclonal antibody to PIGF
Purification:	The antibody was purified by immunogen affinity chromatography.

## Target Details

Target:	PIGF
Alternative Name:	PIGF ( <a href="#">PIGF Products</a> )
Background:	Phosphatidylinositol-glycan biosynthesis class F protein, PIG-F, GPI11 homolog

## Target Details

Gene ID:	5281, 18701
UniProt:	<a href="#">Q07326</a> , <a href="#">O09101</a>
Pathways:	<a href="#">Inositol Metabolic Process</a>


## Application Details

Application Notes:	WB (1:500 - 1:1000), IH (1:100 - 1:200)
Restrictions:	For Research Use only

## Handling


Format:	Liquid
Buffer:	Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.
Expiry Date:	12 months

## Images



### Immunohistochemistry

**Image 1.** Immunohistochemical analysis of PIGF staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Western Blotting

**Image 2.** Western blot analysis of PIGF expression in HEK293T (A), Raw264.7 (B), H9C2 (C) whole cell lysates.