

Datasheet for ABIN7301154
anti-PRKAG2 antibody (N-Term)[Go to Product page](#)

3 Images

Overview

Quantity:	100 µL
Target:	PRKAG2
Binding Specificity:	N-Term
Reactivity:	Human, Monkey
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PRKAG2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF), Immunochromatography (IC)

Product Details

Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human AMPK gamma 2.
Specificity:	Recognizes endogenous levels of AMPK gamma 2 protein.
Characteristics:	Rabbit polyclonal antibody to AMPK gamma 2
Purification:	The antibody was purified by immunogen affinity chromatography.

Target Details

Target:	PRKAG2
Alternative Name:	AMPK gamma 2 (PRKAG2 Products)

Target Details


Background:	5'-AMP-activated protein kinase subunit gamma-2, AMPK gamma2, AMPK subunit gamma-2, H91620p
Gene ID:	51422, 108099
UniProt:	Q9UGJ0 , Q91WG5
Pathways:	AMPK Signaling , Cellular Glucan Metabolic Process , Ribonucleoside Biosynthetic Process , Regulation of Carbohydrate Metabolic Process , Warburg Effect

Application Details

Application Notes:	WB (1:500 - 1:1000), IH (1:100 - 1:200), IF/IC (1:100 - 1:500)
Restrictions:	For Research Use only


Handling

Format:	Liquid
Buffer:	Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.
Expiry Date:	12 months




Immunohistochemistry

Image 1. Immunohistochemical analysis of AMPK gamma 2 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Western Blotting

Image 2. Western blot analysis of AMPK gamma 2 expression in MCF7 (A), HeLa (B) whole cell lysates.



Immunofluorescence

Image 3. Immunofluorescent analysis of AMPK gamma 2 staining in MCF7 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).