



[Go to Product page](#)

Datasheet for ABIN7301268

## anti-SDCCAG8 antibody (C-Term)

### 2 Images

#### Overview

Quantity:	100 µL
Target:	SDCCAG8
Binding Specificity:	C-Term
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SDCCAG8 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC)

#### Product Details

Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human SDCCAG8.
Specificity:	Recognizes endogenous levels of SDCCAG8 protein.
Characteristics:	Rabbit polyclonal antibody to SDCCAG8
Purification:	The antibody was purified by immunogen affinity chromatography.

#### Target Details

Target:	SDCCAG8
Alternative Name:	SDCCAG8 ( <a href="#">SDCCAG8 Products</a> )
Background:	CCCAP, NPHP10, Serologically defined colon cancer antigen 8, Antigen NY-CO-8, Centrosomal

## Target Details

colon cancer autoantigen protein, hCCCAP

Gene ID: 10806, 76816

UniProt: [Q86SQ7](#), [Q80UF4](#)

Pathways: [M Phase](#), [Tube Formation](#)

## Application Details

Application Notes: WB (1:500 - 1:1000), IH (1:100 - 1:200)

Restrictions: For Research Use only

## Handling

Format: Liquid

Buffer: Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.

Preservative: Sodium azide


Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.

Expiry Date: 12 months


## Images



### Immunohistochemistry

**Image 1.** Immunohistochemical analysis of SDCCAG8 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with

haematoxylin and mounted with DPX.



### Western Blotting

**Image 2.** Western blot analysis of SDCCAG8 expression in HEK293T (A), Raw264.7 (B), H9C2 (C) whole cell lysates.