



Datasheet for ABIN7301356  
**anti-TAF13 antibody (Center)**



[Go to Product page](#)

2 Images

Overview

Quantity:	100 µL
Target:	TAF13
Binding Specificity:	Center
Reactivity:	Human, Mouse, Rat, Cow, Zebrafish (Danio rerio)
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TAF13 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC)

Product Details

Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human TAF13.
Specificity:	Recognizes endogenous levels of TAF13 protein.
Characteristics:	Rabbit polyclonal antibody to TAF13
Purification:	The antibody was purified by immunogen affinity chromatography.

Target Details

Target:	TAF13
Alternative Name:	TAF13 ( <a href="#">TAF13 Products</a> )
Background:	TAF2K, TAFII18, Transcription initiation factor TFIID subunit 13, Transcription initiation factor

## Target Details

TFIID 18 kDa subunit, TAF(II)18, TAFII-18, TAFII18

Gene ID: 6884, 99730

UniProt: [Q15543](#), [P61216](#)

## Application Details

Application Notes: WB (1:500 - 1:1000), IH (1:100 - 1:200)

Restrictions: For Research Use only

## Handling

Format: Liquid

Buffer: Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.

Preservative: Sodium azide


Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.


Expiry Date: 12 months

## Images



### Immunohistochemistry

**Image 1.** Immunohistochemical analysis of TAF13 staining in human lung cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



### Western Blotting

**Image 2.** Western blot analysis of TAF13 expression in HEK293T (A), H9C2 (B) whole cell lysates.