antibodies - online.com







anti-TNNI3K antibody (Center)

Images



$C \circ + \circ$	Product	nogo
G0 10	Produci	Luaue

	rv/		

Overview	
Quantity:	100 μL
Target:	TNNI3K
Binding Specificity:	Center
Reactivity:	Human, Mouse, Rat, Cow
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TNNI3K antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC)
Product Details	
Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human CARK.
Specificity:	Recognizes endogenous levels of CARK protein.
Characteristics:	Rabbit polyclonal antibody to CARK
Purification:	The antibody was purified by immunogen affinity chromatography.
Target Details	
Target:	TNNI3K
Alternative Name:	CARK (TNNI3K Products)
Background:	CARK, Serine/threonine-protein kinase TNNI3K, Cardiac ankyrin repeat kinase, Cardiac troponin

Target Details

	I-interacting kinase, TNNI3-interacting kinase
Gene ID:	100526835, 51086, 435766
UniProt:	Q59H18, Q5GIG6, Q7TQP6


Application Details

Application Notes:	WB (1:500 - 1:1000), IH (1:100 - 1:200)
Restrictions:	For Research Use only

Handling


Format:	Liquid
Buffer:	Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.
Expiry Date:	12 months

Images



Immunohistochemistry

Image 1. Immunohistochemical analysis of CARK staining in human heart formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugad compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Western Blotting

Image 2. Western blot analysis of CARK expression in HEK293T (A), Raw264.7 (B), H9C2 (C) whole cell lysates.