

Datasheet for ABIN7301408
anti-TRADD antibody (C-Term)



[Go to Product page](#)

2 Images

Overview

Quantity:	100 µL
Target:	TRADD
Binding Specificity:	C-Term
Reactivity:	Human, Mouse, Rat, Monkey, Dog, Cow, Pig
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TRADD antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC)

Product Details

Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human TRADD.
Specificity:	Recognizes endogenous levels of TRADD protein.
Characteristics:	Rabbit polyclonal antibody to TRADD
Purification:	The antibody was purified by immunogen affinity chromatography.

Target Details

Target:	TRADD
Alternative Name:	TRADD (TRADD Products)
Background:	Tumor necrosis factor receptor type 1-associated DEATH domain protein, TNFR1-associated

Target Details


	DEATH domain protein, TNFRSF1A-associated via death domain
Gene ID:	8717, 71609
UniProt:	Q15628 , Q3U0V2
Pathways:	NF-kappaB Signaling , Apoptosis , Caspase Cascade in Apoptosis , Positive Regulation of Endopeptidase Activity , Hepatitis C

Application Details

Application Notes:	WB (1:500 - 1:1000), IH (1:100 - 1:200)
Restrictions:	For Research Use only


Handling

Format:	Liquid
Buffer:	Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.
Expiry Date:	12 months



Immunohistochemistry

Image 1. Immunohistochemical analysis of TRADD staining in human colon cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Western Blotting

Image 2. Western blot analysis of TRADD expression in K562 (A), mouse heart (B), rat liver (C) whole cell lysates.