antibodies -online.com





anti-UBA5 antibody (C-Term)

2 Images



Go to Product page

\sim			
()	N/Ar	\sqrt{i}	

Overview	
Quantity:	100 μL
Target:	UBA5
Binding Specificity:	C-Term
Reactivity:	Human, Monkey
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This UBA5 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC)
Product Details	
Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human UBA5.
Specificity:	Recognizes endogenous levels of UBA5 protein.
Characteristics:	Rabbit polyclonal antibody to UBA5
Purification:	The antibody was purified by immunogen affinity chromatography.
Target Details	
Target:	UBA5
Alternative Name:	UBA5 (UBA5 Products)
Background:	UBE1DC1, Ubiquitin-like modifier-activating enzyme 5, Ubiquitin-activating enzyme 5, ThiFP1,

Target Details

	UFM1-activating enzyme, Ubiquitin-activating enzyme E1 domain-containing protein 1
Gene ID:	79876, 66663, 300968
UniProt:	Q9GZZ9, Q8VE47, Q5M7A4


Application Details

Application Notes:	WB (1:500 - 1:1000), IH (1:100 - 1:200)
Restrictions:	For Research Use only

Handling


Format:	Liquid
Buffer:	Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.
Expiry Date:	12 months

Images



Western Blotting

Image 1. Western blot analysis of UBA5 expression in HEK293T (A) whole cell lysates.



Immunohistochemistry

Image 2. Immunohistochemical analysis of UBA5 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugad compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.