

Datasheet for ABIN7301428
anti-Ubiquilin 3 antibody (Center)

2 Images

[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	Ubiquilin 3 (UBQLN3)
Binding Specificity:	Center
Reactivity:	Human, Rat, Cow, Monkey
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Ubiquilin 3 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC)

Product Details

Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human Ubiquilin 3.
Specificity:	Recognizes endogenous levels of Ubiquilin 3 protein.
Characteristics:	Rabbit polyclonal antibody to Ubiquilin 3
Purification:	The antibody was purified by immunogen affinity chromatography.

Target Details

Target:	Ubiquilin 3 (UBQLN3)
Alternative Name:	Ubiquilin 3 (UBQLN3 Products)
Background:	Ubiquilin-3

Target Details

Gene ID: 50613, 244178

UniProt: [Q9H347](#), [Q8C5U9](#)

Application Details

Application Notes: WB (1:500 - 1:1000), IH (1:100 - 1:200)

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.

Preservative: Sodium azide


Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.


Expiry Date: 12 months

Images



Immunohistochemistry

Image 1. Immunohistochemical analysis of Ubiquilin 3 staining in human testis formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Western Blotting

Image 2. Western blot analysis of Ubiquilin 3 expression in HepG2 (A), rat brain (B), rat liver (C) whole cell lysates.