



Datasheet for ABIN7301458
anti-ZNF232 antibody (Center)



[Go to Product page](#)

2 Images

Overview

| | |
|----------------------|---|
| Quantity: | 100 µL |
| Target: | ZNF232 |
| Binding Specificity: | Center |
| Reactivity: | Human, Mouse, Rat, Monkey |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This ZNF232 antibody is un-conjugated |
| Application: | Western Blotting (WB), Immunohistochemistry (IHC) |

Product Details

| | |
|------------------|--|
| Immunogen: | KLH-conjugated synthetic peptide encompassing a sequence within the center region of human ZNF232. |
| Specificity: | Recognizes endogenous levels of ZNF232 protein. |
| Characteristics: | Rabbit polyclonal antibody to ZNF232 |
| Purification: | The antibody was purified by immunogen affinity chromatography. |

Target Details

| | |
|-------------------|---|
| Target: | ZNF232 |
| Alternative Name: | ZNF232 (ZNF232 Products) |
| Background: | ZSCAN11, Zinc finger protein 232, Zinc finger and SCAN domain-containing protein 11 |

Target Details

Gene ID: 7775

UniProt: [Q9UNY5](#)

Application Details

Application Notes: WB (1:500 - 1:1000), IH (1:100 - 1:200)

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.

Preservative: Sodium azide


Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.


Expiry Date: 12 months

Images



Immunohistochemistry

Image 1. Immunohistochemical analysis of ZNF232 staining in human testis formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Western Blotting

Image 2. Western blot analysis of ZNF232 expression in HEK293T (A), NS-1 (B), H9C2 (C) whole cell lysates.