



[Go to Product page](#)

Datasheet for ABIN7301462
anti-ZNF76 antibody (Center)

2 Images

Overview

Quantity:	100 µL
Target:	ZNF76
Binding Specificity:	Center
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ZNF76 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC)

Product Details

Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human ZNF76.
Specificity:	Recognizes endogenous levels of ZNF76 protein.
Characteristics:	Rabbit polyclonal antibody to ZNF76
Purification:	The antibody was purified by immunogen affinity chromatography.

Target Details

Target:	ZNF76
Alternative Name:	ZNF76 (ZNF76 Products)
Background:	D6S229E, ZNF523, Zinc finger protein 76, Zinc finger protein 523

Target Details

Gene ID: 7629, 224656, 361809

UniProt: [P36508](#), [Q8BMU0](#), [B4F7E9](#)

Application Details

Application Notes: WB (1:500 - 1:1000), IH (1:100 - 1:200)

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Liquid in 0.42 % Potassium phosphate, 0.87 % Sodium chloride, pH 7.3, 30 % glycerol, and 0.01 % sodium azide.

Preservative: Sodium azide


Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.


Expiry Date: 12 months

Images



Western Blotting

Image 1. Western blot analysis of ZNF76 expression in HEK293T (A), Raw264.7 (B), PC12 (C) whole cell lysates.



Immunohistochemistry

Image 2. Immunohistochemical analysis of ZNF76 staining in human testis formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.