

Datasheet for ABIN730267 anti-Endothelin 1 antibody (AA 53-73) (HRP)



Overview

| 0.10.1.011 | |
|-----------------------|---|
| Quantity: | 100 μL |
| Target: | Endothelin 1 (EDN1) |
| Binding Specificity: | AA 53-73 |
| Reactivity: | Human, Rat, Pig, Mouse |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This Endothelin 1 antibody is conjugated to HRP |
| Application: | Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)) |
| Product Details | |
| Immunogen: | KLH conjugated synthetic peptide derived from human human Endothelin-1 |
| Isotype: | IgG |
| Cross-Reactivity: | Human, Mouse, Pig, Rat |
| Predicted Reactivity: | Dog,Cow,Sheep,Horse,Guinea Pig |
| Purification: | Purified by Protein A. |
| Target Details | |
| Target: | Endothelin 1 (EDN1) |
| Alternative Name: | Endothelin-1 (EDN1 Products) |

Target Details

| Background: | Synonyms: Endothelin-1, EDN 1, EDN1, EDN1 protein, Endothelin1, ET 1,ET1, |
|---------------------|---|
| | Background: Endothelins (ET) show potent constrictor activity in vascular and non-vascular |
| | smooth muscle. This family of 21-amino acid peptides exists in at least three isoforms - ET-1, |
| | ET-2, and ET-3, and is produced in endothelial and epithelial cells. ET's can mediate biological |
| | effects in cells and tissues, and have been shown to bind to an ET receptor in the lung, kidney, |
| | heart, and liver. Endothelin 1 is expressed in lung, placental stem villi vessels and in cultured |
| | placental vascular smooth muscle cells. |
| Gene ID: | 1906 |
| Pathways: | Hormone Transport, Negative Regulation of Hormone Secretion, Regulation of Systemic Arteria |
| | Blood Pressure by Hormones, cAMP Metabolic Process, Regulation of Muscle Cell |
| | Differentiation, Regulation of G-Protein Coupled Receptor Protein Signaling, Regulation of Cell |
| | Size |
| Application Details | |
| Application Notes: | WB 1:300-5000 |
| | IHC-P 1:200-400 |
| | IHC-F 1:100-500 |
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Liquid |
| Concentration: | 1 μg/μL |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and |
| | 50 % Glycerol. |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be |
| | handled by trained staff only. |
| Storage: | -20 °C |
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. |
| Expiry Date: | 12 months |
| | |