

Datasheet for ABIN7302738

anti-ZNF265 antibody**3** Images[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	ZNF265 (Zranb2)
Reactivity:	Human, Mouse, Rat, Dog, Rabbit, Pig, Chicken, Monkey
Host:	Rabbit
Clonality:	Polyclonal
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF), Immunochromatography (IC)

Product Details

Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human ZNF265.
Specificity:	Recognizes endogenous levels of ZNF265 protein.
Characteristics:	Rabbit polyclonal antibody to ZNF265
Purification:	The antibody was purified by affinity chromatography.

Target Details

Target:	ZNF265 (Zranb2)
Alternative Name:	ZNF265 (Zranb2 Products)
Background:	ZIS, ZNF265, Zinc finger Ran-binding domain-containing protein 2, Zinc finger protein 265, Zinc finger, splicing
Gene ID:	9406, 53861, 58821

Target Details

UniProt: [O95218](#), [Q9R020](#), [O35986](#)

Application Details

Application Notes: WB (1:500 - 1:1000), IH (1:100 - 1:200), IF/IC (1:100 - 1:500)

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Liquid in PBS, pH 7.3, 0.2 % BSA, and 0.02 % sodium azide.

Preservative: Sodium azide


Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.


Expiry Date: 12 months

Images




Immunofluorescence

Image 1. Immunofluorescent analysis of ZNF265 staining in HepG2 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody



Immunohistochemistry

Image 2. Immunohistochemical analysis of ZNF265 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated w



Western Blotting

Image 3. Western blot analysis of ZNF265 expression in HepG2 (A), MCF7 (B) whole cell lysates.