antibodies -online.com





anti-NCBP2 antibody (N-Term)

3 Images



Go to Product page

\sim				
	11/6	٦r١	/10	۱۸.

Alternative Name:

Quantity:	100 μL	
Target:	NCBP2	
Binding Specificity:	N-Term	
Reactivity:	Human, Mouse, Rat, Cow, Monkey	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This NCBP2 antibody is un-conjugated	
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF),	
	Immunochromatography (IC)	
Product Details		
Jadot Dotallo		
Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of	
	KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human CBP20.	
Immunogen:	human CBP20.	
Immunogen: Specificity:	human CBP20. Recognizes endogenous levels of CBP20 protein.	
Immunogen: Specificity: Characteristics:	human CBP20. Recognizes endogenous levels of CBP20 protein. Rabbit polyclonal antibody to CBP20	


CBP20 (NCBP2 Products)

Target Details

Expiry Date:


12 months

rai got botano		
Background:	CBP20, Nuclear cap-binding protein subunit 2, 20 kDa nuclear cap-binding protein, Cell proliferation-inducing gene 55 protein, NCBP 20 kDa subunit, CBP20, NCBP-interacting protein 1, NIP1	
Gene ID:	22916, 68092, 689116	
UniProt:	P52298, Q9CQ49, B1WC40	
Pathways:	Ribonucleoprotein Complex Subunit Organization, Methionine Biosynthetic Process	
Application Details		
Application Notes:	WB (1:500 - 1:1000), IH (1:100 - 1:200), IF/IC (1:100 - 1:500)	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Buffer:	Liquid in PBS, pH 7.3, 0.2 % BSA, and 0.02 % sodium azide.	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.	
F : D :	40	




Immunofluorescence

Image 1. Immunofluorescent analysis of CBP20 staining in Hela cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody



Immunohistochemistry

Image 2. Immunohistochemical analysis of CBP20 staining in human prostate cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated



Western Blotting

Image 3. Western blot analysis of CBP20 expression in Hela (A), NIH3T3 (B) whole cell lysates.