

Datasheet for ABIN7302772

anti-DPF2 antibody**3** Images[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	DPF2
Reactivity:	Human, Mouse, Rat, Cow
Host:	Rabbit
Clonality:	Polyclonal
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF), Immunochromatography (IC)

Product Details

Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human DPF2.
Specificity:	Recognizes endogenous levels of DPF2 protein.
Characteristics:	Rabbit polyclonal antibody to DPF2
Purification:	The antibody was purified by affinity chromatography.

Target Details

Target:	DPF2
Alternative Name:	DPF2 (DPF2 Products)
Background:	BAF45D, REQ, UBID4, Zinc finger protein ubi-d4, Apoptosis response zinc finger protein, BRG1-associated factor 45D, BAF45D, D4, zinc and double PHD fingers family 2, Protein requiem
Gene ID:	5977, 19708

Target Details

UniProt: [Q92785](#), [Q61103](#)

Application Details

Application Notes: WB (1:500 - 1:1000), IH (1:100 - 1:200), IF/IC (1:100 - 1:500)

Restrictions: For Research Use only

Handling

Format: Liquid

Buffer: Liquid in PBS, pH 7.3, 0.2 % BSA, and 0.02 % sodium azide.

Preservative: Sodium azide


Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.


Expiry Date: 12 months

Images




Immunofluorescence

Image 1. Immunofluorescent analysis of DPF2 staining in Jurkat cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody



Immunohistochemistry

Image 2. Immunohistochemical analysis of DPF2 staining in human lung cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with



Western Blotting

Image 3. Western blot analysis of DPF2 expression in Jurkat (A), THP1 (B), K562 (C) whole cell lysates.