

Datasheet for ABIN7302918

anti-Tyrosine Hydroxylase antibody (N-Term, pSer8)



[Go to Product page](#)

2 Images

Overview

Quantity:	100 µL
Target:	Tyrosine Hydroxylase (TH)
Binding Specificity:	N-Term, pSer8
Reactivity:	Human, Rat, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Tyrosine Hydroxylase antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP)

Product Details

Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human Tyrosine Hydroxylase.
Specificity:	Recognizes endogenous levels of Tyrosine Hydroxylase (pS8) protein.
Characteristics:	Rabbit polyclonal antibody to Tyrosine Hydroxylase (pS8)
Purification:	The antibody was purified by affinity chromatography.

Target Details

Target:	Tyrosine Hydroxylase (TH)
Alternative Name:	Tyrosine Hydroxylase (TH Products)
Background:	TH, TYH, Tyrosine 3-monooxygenase, Tyrosine 3-hydroxylase, TH

Target Details

Gene ID:	25085
UniProt:	P07101 , P04177
Pathways:	Dopaminergic Neurogenesis , Response to Water Deprivation , Sensory Perception of Sound , Carbohydrate Homeostasis , Feeding Behaviour


Application Details

Application Notes:	WB (1:500 - 1:1000), IH (1:100 - 1:200), IP (1:10 - 1:100)
Restrictions:	For Research Use only

Handling


Format:	Liquid
Buffer:	Liquid in PBS, pH 7.3, 0.2 % BSA, and 0.02 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.
Expiry Date:	12 months

Images



Immunohistochemistry

Image 1. Immunohistochemical analysis of Tyrosine Hydroxylase (pS8) staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then



Western Blotting

Image 2. Western blot analysis of Tyrosine Hydroxylase (pS8) expression in HEK293T (A), HepG2 (B), NIH3T3 (C) whole cell lysates.