

Datasheet for ABIN7303102

**anti-ZNF668 antibody****3** Images[Go to Product page](#)

## Overview

Quantity:	100 µL
Target:	ZNF668
Reactivity:	Human, Mouse, Rat, Cow, Monkey
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ZNF668 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF), Immunochromatography (IC)

## Product Details

Immunogen:	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human ZNF668.
Specificity:	Recognizes endogenous levels of ZNF668 protein.
Characteristics:	Rabbit polyclonal antibody to ZNF668
Purification:	The antibody was purified by affinity chromatography.

## Target Details

Target:	ZNF668
Alternative Name:	ZNF668 ( <a href="#">ZNF668 Products</a> )
Background:	Zinc finger protein 668

## Target Details

Gene ID: 79759, 244219

UniProt: [Q96K58](#), [Q8K2R5](#)

## Application Details

Application Notes: WB (1:500 - 1:1000), IH (1:100 - 1:200), IF/IC (1:100 - 1:500)

Restrictions: For Research Use only

## Handling

Format: Liquid

Buffer: Liquid in PBS, pH 7.3, 0.2 % BSA, and 0.02 % sodium azide.

Preservative: Sodium azide


Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.


Expiry Date: 12 months

## Images




### Immunofluorescence

**Image 1.** Immunofluorescent analysis of ZNF668 staining in HepG2 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody



### Immunohistochemistry

**Image 2.** Immunohistochemical analysis of ZNF668 staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the



### Western Blotting

**Image 3.** Western blot analysis of ZNF668 expression in HepG2 (A) whole cell lysates.